

Time series analysis

Lecture 8. Regressive dynamic models

Dr. Khamidov Obidjon

Violation of the Assumptions of the CLRM

- Recall that we assumed of the CLRM disturbance terms:
 1. $E(u_t) = 0$
 2. $\text{Var}(u_t) = \sigma^2 < \infty$
 3. $\text{Cov}(u_i, u_j) = 0$
 4. The X matrix is non-stochastic or fixed in repeated samples
 5. $u_t \sim N(0, \sigma^2)$

Investigating Violations of the Assumptions of the CLRM

- We will now study these assumptions further, and in particular look at:
 - How we test for violations
 - Causes
 - Consequences
 - in general we could encounter any combination of 3 problems:
 - the coefficient estimates are wrong
 - the associated standard errors are wrong
 - the distribution that we assumed for the test statistics will be inappropriate
 - Solutions
 - the assumptions are no longer violated
 - we work around the problem so that we use alternative techniques which are still valid

Statistical Distributions for Diagnostic Tests


- Often, an F - and a χ^2 - version of the test are available.
- The F -test version involves estimating a restricted and an unrestricted version of a test regression and comparing the RSS .
- The χ^2 - version is sometimes called an “LM” test, and only has one degree of freedom parameter: the number of restrictions being tested, m .
- Asymptotically, the 2 tests are equivalent since the χ^2 is a special case of the F -distribution:
$$\frac{\chi^2(m)}{m} \rightarrow F(m, T - k) \text{ as } T - k \rightarrow \infty$$
- For small samples, the F -version is preferable.

Assumption 1: $E(u_t) = 0$

- Assumption that the mean of the disturbances is zero.
- For all diagnostic tests, we cannot observe the disturbances and so perform the tests of the residuals.
- The mean of the residuals will always be zero provided that there is a constant term in the regression.

Assumption 2: $\text{Var}(u_t) = \sigma^2 < \infty$

- We have so far assumed that the variance of the errors is constant, σ^2 - this is known as homoscedasticity.
- If the errors do not have a constant variance, we say that they are heteroscedastic e.g. say we estimate a regression and calculate the residuals, \hat{u}_t .



Detection of Heteroscedasticity: The GQ Test

- Graphical methods
- Formal tests: There are many of them: we will discuss Goldfeld-Quandt test and White's test

The Goldfeld-Quandt (GQ) test is carried out as follows.

1. Split the total sample of length T into two sub-samples of length T_1 and T_2 . The regression model is estimated on each sub-sample and the two residual variances are calculated.
2. The null hypothesis is that the variances of the disturbances are equal,

$$H_0: \sigma_1^2 = \sigma_2^2$$

The GQ Test (Cont'd)

3. The test statistic, denoted GQ , is simply the ratio of the two residual variances where the larger of the two variances must be placed in the numerator.

$$GQ = \frac{s_1^2}{s_2^2}$$

4. The test statistic is distributed as an $F(T_1-k, T_2-k)$ under the null of homoscedasticity.
5. A problem with the test is that the choice of where to split the sample is that usually arbitrary and may crucially affect the outcome of the test.

Detection of Heteroscedasticity using White's Test

- White's general test for heteroscedasticity is one of the best approaches because it makes few assumptions about the form of the heteroscedasticity.
- The test is carried out as follows:

1. Assume that the regression we carried out is as follows

$$y_t = \beta_1 + \beta_2 x_{2t} + \beta_3 x_{3t} + u_t$$

And we want to test $\text{Var}(u_t) = \sigma^2$. We estimate the model, obtaining the residuals,

$$\hat{u}_t$$

2. Then run the auxiliary regression

$$\hat{u}_t^2 = \alpha_1 + \alpha_2 x_{2t} + \alpha_3 x_{3t} + \alpha_4 x_{2t}^2 + \alpha_5 x_{3t}^2 + \alpha_6 x_{2t} x_{3t} + v_t$$

Performing White's Test for Heteroscedasticity

3. Obtain R^2 from the auxiliary regression and multiply it by the number of observations, T . It can be shown that

$$T R^2 \sim \chi^2 (m)$$

where m is the number of regressors in the auxiliary regression excluding the constant term.

4. If the χ^2 test statistic from step 3 is greater than the corresponding value from the statistical table then reject the null hypothesis that the disturbances are homoscedastic.

Consequences of Using OLS in the Presence of Heteroscedasticity

- OLS estimation still gives unbiased coefficient estimates, but they are no longer BLUE.
- This implies that if we still use OLS in the presence of heteroscedasticity, our standard errors could be inappropriate and hence any inferences we make could be misleading.
- Whether the standard errors calculated using the usual formulae are too big or too small will depend upon the form of the heteroscedasticity.

How Do we Deal with Heteroscedasticity?

- If the form (i.e. the cause) of the heteroscedasticity is known, then we can use an estimation method which takes this into account (called generalised least squares, GLS).
- A simple illustration of GLS is as follows: Suppose that the error variance is related to another variable z_t by

$$\text{var}(u_t) = \sigma^2 z_t^2$$

- To remove the heteroscedasticity, divide the regression equation by z_t

$$\frac{y_t}{z_t} = \beta_1 \frac{1}{z_t} + \beta_2 \frac{x_{2t}}{z_t} + \beta_3 \frac{x_{3t}}{z_t} + v_t$$

where $v_t = \frac{u_t}{z_t}$ is an error term.

- Now $\text{var}(v_t) = \text{var}\left(\frac{u_t}{z_t}\right) = \frac{\text{var}(u_t)}{z_t^2} = \frac{\sigma^2 z_t^2}{z_t^2} = \sigma^2$ for known z_t .

Other Approaches to Dealing with Heteroscedasticity

- So the disturbances from the new regression equation will be homoscedastic.
- Other solutions include:
 1. Transforming the variables into logs or reducing by some other measure of “size”.
 2. Use White’s heteroscedasticity consistent standard error estimates.

The effect of using White’s correction is that in general the standard errors for the slope coefficients are increased relative to the usual OLS standard errors.

This makes us more “conservative” in hypothesis testing, so that we would need more evidence against the null hypothesis before we would reject it.


Background – The Concept of a Lagged Value

t	y_t	y_{t-1}	Δy_t
1989M09	0.8	-	-
1989M10	1.3	0.8	$1.3-0.8=0.5$
1989M11	-0.9	1.3	$-0.9-1.3=-2.2$
1989M12	0.2	-0.9	$0.2--0.9=1.1$
1990M01	-1.7	0.2	$-1.7-0.2=-1.9$
1990M02	2.3	-1.7	$2.3--1.7=4.0$
1990M03	0.1	2.3	$0.1-2.3=-2.2$
1990M04	0.0	0.1	$0.0-0.1=-0.1$
.	.	.	.
.	.	.	.
.	.	.	.

Autocorrelation



- We assumed of the CLRM's errors that $\text{Cov}(u_i, u_j) = 0$ for $i \neq j$, i.e. This is essentially the same as saying there is no pattern in the errors.
- Obviously we never have the actual u 's, so we use their sample counterpart, the residuals (the \hat{u}_t 's).
- If there are patterns in the residuals from a model, we say that they are autocorrelated.
- Some stereotypical patterns we may find in the residuals are given on the next 3 slides.

Positive Autocorrelation





Positive Autocorrelation is indicated by a cyclical residual plot over time.

Negative Autocorrelation



Negative autocorrelation is indicated by an alternating pattern where the residuals cross the time axis more frequently than if they were distributed randomly

No pattern in residuals – No autocorrelation



No pattern in residuals at all: this is what we would like to see

Detecting Autocorrelation: The Durbin-Watson Test

The Durbin-Watson (DW) is a test for first order autocorrelation - i.e. it assumes that the relationship is between an error and the previous one

$$u_t = \rho u_{t-1} + v_t \quad (1)$$

where $v_t \sim N(0, \sigma_v^2)$.

- The DW test statistic actually tests

$$H_0 : \rho=0 \text{ and } H_1 : \rho \neq 0$$

- The test statistic is calculated by

$$DW = \frac{\sum_{t=2}^T (\hat{u}_t - \hat{u}_{t-1})^2}{\sum_{t=2}^T \hat{u}_t^2}$$

The Durbin-Watson Test: Critical Values

- We can also write


$$DW \approx 2(1 - \hat{\rho}) \quad (2)$$

where $\hat{\rho}$ is the estimated correlation coefficient. Since $\hat{\rho}$ is a correlation, it implies that $-1 \leq \hat{\rho} \leq 1$

- Rearranging for DW from (2) would give $0 \leq DW \leq 4$.
- If $\hat{\rho} = 0$, $DW = 2$. So roughly speaking, do not reject the null hypothesis if DW is near 2 \rightarrow i.e. there is little evidence of autocorrelation
- Unfortunately, DW has 2 critical values, an upper critical value (d_u) and a lower critical value (d_L), and there is also an intermediate region where we can neither reject nor not reject H_0 .

The Durbin-Watson Test: Interpreting the Results

Reject H_0 : positive autocorrelation Inconclusive Do not reject H_0 : No evidence of autocorrelation Inconclusive Reject H_0 : negative autocorrelation



Conditions which Must be Fulfilled for DW to be a Valid Test

1. Constant term in regression
2. Regressors are non-stochastic
3. No lags of dependent variable

Another Test for Autocorrelation: The Breusch-Godfrey Test

- It is a more general test for r^{th} order autocorrelation:

$$u_t = \rho_1 u_{t-1} + \rho_2 u_{t-2} + \rho_3 u_{t-3} + \dots + \rho_r u_{t-r} + v_t \quad , \quad v_t \sim N(0, \sigma_v^2)$$

- The null and alternative hypotheses are:

$$H_0 : \rho_1 = 0 \text{ and } \rho_2 = 0 \text{ and } \dots \text{ and } \rho_r = 0$$

$$H_1 : \rho_1 \neq 0 \text{ or } \rho_2 \neq 0 \text{ or } \dots \text{ or } \rho_r \neq 0$$

- The test is carried out as follows:

1. Estimate the linear regression using OLS and obtain the residuals, \hat{u}_t .

2. Regress \hat{u}_t on all of the regressors from stage 1 (the x 's) plus $\hat{u}_{t-1}, \hat{u}_{t-2}, \dots, \hat{u}_{t-r}$
Obtain R^2 from this regression.

3. It can be shown that $(T-r)R^2 \sim \chi^2(r)$

- If the test statistic exceeds the critical value from the statistical tables, reject the null hypothesis of no autocorrelation.

Consequences of Ignoring Autocorrelation if it is Present

- The coefficient estimates derived using OLS are still unbiased, but they are inefficient, i.e. they are not BLUE, even in large sample sizes.
- Thus, if the standard error estimates are inappropriate, there exists the possibility that we could make the wrong inferences.
- R^2 is likely to be inflated relative to its “correct” value for positively correlated residuals.

“Remedies” for Autocorrelation

- If the form of the autocorrelation is known, we could use a GLS procedure – i.e. an approach that allows for autocorrelated residuals e.g., Cochrane-Orcutt.
- But such procedures that “correct” for autocorrelation require assumptions about the form of the autocorrelation.
- If these assumptions are invalid, the cure would be more dangerous than the disease! - see Hendry and Mizon (1978).
- However, it is unlikely to be the case that the form of the autocorrelation is known, and a more “modern” view is that residual autocorrelation presents an opportunity to modify the regression.

Dynamic Models

- All of the models we have considered so far have been static, e.g.

$$y_t = \beta_1 + \beta_2 x_{2t} + \dots + \beta_k x_{kt} + u_t$$

- But we can easily extend this analysis to the case where the current value of y_t depends on previous values of y or one of the x 's, e.g.

$$y_t = \beta_1 + \beta_2 x_{2t} + \dots + \beta_k x_{kt} + \gamma_1 y_{t-1} + \gamma_2 x_{2t-1} + \dots + \gamma_k x_{kt-1} + u_t$$

- We could extend the model even further by adding extra lags, e.g.

$$x_{2t-2}, y_{t-3}.$$

Why Might we Want/Need To Include Lags in a Regression?

- Inertia of the dependent variable
- Over-reactions
- Measuring time series as overlapping moving averages
- However, other problems with the regression could cause the null hypothesis of no autocorrelation to be rejected:
 - Omission of relevant variables, which are themselves autocorrelated.
 - If we have committed a “misspecification” error by using an inappropriate functional form.
 - Autocorrelation resulting from unparameterised seasonality.

Models in First Difference Form

- Another way to sometimes deal with the problem of autocorrelation is to switch to a model in first differences.
- Denote the first difference of y_t , i.e. $y_t - y_{t-1}$ as Δy_t ; similarly for the x -variables, $\Delta x_{2t} = x_{2t} - x_{2t-1}$ etc.

- The model would now be

$$\Delta y_t = \beta_1 + \beta_2 \Delta x_{2t} + \dots + \beta_k \Delta x_{kt} + u_t$$

- Sometimes the change in y is purported to depend on previous values of y or x_t as well as changes in x :

$$\Delta y_t = \beta_1 + \beta_2 \Delta x_{2t} + \beta_3 x_{2t-1} + \beta_4 y_{t-1} + u_t$$

The Long Run Static Equilibrium Solution

- One interesting property of a dynamic model is its long run or static equilibrium solution.
- “Equilibrium” implies that the variables have reached some steady state and are no longer changing, i.e. if y and x are in equilibrium, we can say

$$y_t = y_{t+1} = \dots = y \text{ and } x_t = x_{t+1} = \dots = x$$

Consequently, $\Delta y_t = y_t - y_{t-1} = y - y = 0$ etc.

- So the way to obtain a long run static solution is:
 1. Remove all time subscripts from variables
 2. Set error terms equal to their expected values, $E(u_t)=0$
 3. Remove first difference terms altogether
 4. Gather terms in x together and gather terms in y together.
- These steps can be undertaken in any order

The Long Run Static Equilibrium Solution: An Example

If our model is

$$\Delta y_t = \beta_1 + \beta_2 \Delta x_{2t} + \beta_3 x_{2t-1} + \beta_4 y_{t-1} + u_t$$

then the static solution would be given by

$$0 = \beta_1 + \beta_3 x_{2t-1} + \beta_4 y_{t-1}$$

$$\beta_4 y_{t-1} = -\beta_1 - \beta_3 x_{2t-1}$$

$$y = \frac{-\beta_1}{\beta_4} - \frac{\beta_3}{\beta_4} x_2$$

Problems with Adding Lagged Regressors to “Cure” Autocorrelation

- Inclusion of lagged values of the dependent variable violates the assumption that the RHS variables are non-stochastic.
- What does an equation with a large number of lags actually mean?
- Note that if there is still autocorrelation in the residuals of a model including lags, then the OLS estimators will not even be consistent.

Multicollinearity

- This problem occurs when the explanatory variables are very highly correlated with each other.
- Perfect multicollinearity
Cannot estimate all the coefficients
 - e.g. suppose $x_3 = 2x_2$
 - and the model is $y_t = \beta_1 + \beta_2 x_{2t} + \beta_3 x_{3t} + \beta_4 x_{4t} + u_t$
- Problems if Near Multicollinearity is Present but Ignored
 - R^2 will be high but the individual coefficients will have high standard errors.
 - The regression becomes very sensitive to small changes in the specification.
 - Thus confidence intervals for the parameters will be very wide, and significance tests might therefore give inappropriate conclusions.

Measuring Multicollinearity

- The easiest way to measure the extent of multicollinearity is simply to look at the matrix of correlations between the individual variables. e.g.

Corr	x_2	x_3	x_4
x_2	-	0.2	<u>0.8</u>
x_3	0.2	-	0.3
x_4	<u>0.8</u>	0.3	-

- But another problem: if 3 or more variables are linear
- e.g. $x_{2t} + x_{3t} = x_{4t}$
- Note that high correlation between y and one of the x 's is not multicollinearity.

Solutions to the Problem of Multicollinearity

- “Traditional” approaches, such as ridge regression or principal components. But these usually bring more problems than they solve.
- Some econometricians argue that if the model is otherwise OK, just ignore it
- The easiest ways to “cure” the problems are
 - drop one of the collinear variables
 - transform the highly correlated variables into a ratio
 - go out and collect more data e.g.
 - a longer run of data
 - switch to a higher frequency

Adopting the Wrong Functional Form

- We have previously assumed that the appropriate functional form is linear.
- This may not always be true.
- We can formally test this using Ramsey's RESET test, which is a general test for mis-specification of functional form.

- Essentially the method works by adding higher order terms of the fitted values (e.g. \hat{y}_t^2, \hat{y}_t^3 etc.) into an auxiliary regression:

Regress \hat{u}_t on powers of the fitted values:

$$\hat{u}_t = \beta_0 + \beta_1 \hat{y}_t^2 + \beta_2 \hat{y}_t^3 + \dots + \beta_{p-1} \hat{y}_t^p + v_t$$

Obtain R^2 from this regression. The test statistic is given by TR^2 and is distributed as a $\chi^2(p-1)$.

- So if the value of the test statistic is greater than a $\chi^2(p-1)$ then reject the null hypothesis that the functional form was correct.

But what do we do if this is the case?

- The RESET test gives us no guide as to what a better specification might be.

- One possible cause of rejection of the test is if the true model is

$$y_t = \beta_1 + \beta_2 x_{2t} + \beta_3 x_{2t}^2 + \beta_4 x_{2t}^3 + u_t$$

In this case the remedy is obvious.

- Another possibility is to transform the data into logarithms. This will linearise many previously multiplicative models into additive ones:

$$y_t = Ax_t^\beta e^{u_t} \Leftrightarrow \ln y_t = \alpha + \beta \ln x_t + u_t$$


Testing the Normality Assumption

- Why did we need to assume normality for hypothesis testing?


Testing for Departures from Normality

- *The Bera Jarque normality test*
- A normal distribution is not skewed and is defined to have a coefficient of kurtosis of 3.
- The kurtosis of the normal distribution is 3 so its excess kurtosis (b_2-3) is zero.
- Skewness and kurtosis are the (standardised) third and fourth moments of a distribution.

Normal versus Skewed Distributions




A normal distribution



A skewed distribution

Leptokurtic versus Normal Distribution



Testing for Normality

- Bera and Jarque formalise this by testing the residuals for normality by testing whether the coefficient of skewness and the coefficient of excess kurtosis are jointly zero.
- It can be proved that the coefficients of skewness and kurtosis can be expressed respectively as:

$$b_1 = \frac{E[u^3]}{(\sigma^2)^{3/2}} \quad \text{and} \quad b_2 = \frac{E[u^4]}{(\sigma^2)^2}$$

- The Bera Jarque test statistic is given by


$$W = T \left[\frac{b_1^2}{6} + \frac{(b_2 - 3)^2}{24} \right] \sim \chi^2(2)$$

- We estimate b_1 and b_2 using the residuals from the OLS regression, \hat{u} .

What do we do if we find evidence of Non-Normality?

- It is not obvious what we should do!
- Could use a method which does not assume normality, but difficult and what are its properties?
- Often the case that one or two very extreme residuals causes us to reject the normality assumption.
- An alternative is to use dummy variables.
e.g. say we estimate a monthly model of asset returns from 1980-1990, and we plot the residuals, and find a particularly large outlier for October 1987:

What do we do if we find evidence of Non-Normality? (cont'd)



- Create a new variable:
 $D87M10_t = 1$ during October 1987 and zero otherwise.
This effectively knocks out that observation. But we need a theoretical reason for adding dummy variables.

Omission of an Important Variable or Inclusion of an Irrelevant Variable

Omission of an Important Variable

- Consequence: The estimated coefficients on all the other variables will be biased and inconsistent unless the excluded variable is uncorrelated with all the included variables.
- Even if this condition is satisfied, the estimate of the coefficient on the constant term will be biased.
- The standard errors will also be biased.

Inclusion of an Irrelevant Variable

- Coefficient estimates will still be consistent and unbiased, but the estimators will be inefficient.

Parameter Stability Tests

- So far, we have estimated regressions such as $y_t = \beta_1 + \beta_2 x_{2t} + \beta_3 x_{3t} + u_t$
- We have implicitly assumed that the parameters (β_1 , β_2 and β_3) are constant for the entire sample period.
- We can test this implicit assumption using parameter stability tests. The idea is essentially to split the data into sub-periods and then to estimate up to three models, for each of the sub-parts and for all the data and then to “compare” the *RSS* of the models.
- There are two types of test we can look at:
 - Chow test (analysis of variance test)
 - Predictive failure tests

The Chow Test

- The steps involved are:
 1. Split the data into two sub-periods. Estimate the regression over the whole period and then for the two sub-periods separately (3 regressions). Obtain the RSS for each regression.
 2. The restricted regression is now the regression for the whole period while the “unrestricted regression” comes in two parts: for each of the sub-samples.

We can thus form an F-test which is the difference between the *RSS*'s.

The statistic is
$$\frac{RSS - (RSS_1 + RSS_2)}{RSS_1 + RSS_2} \times \frac{T - 2k}{k}$$

The Chow Test (cont'd)

where:

RSS = RSS for whole sample

RSS_1 = RSS for sub-sample 1

RSS_2 = RSS for sub-sample 2

T = number of observations

$2k$ = number of regressors in the “unrestricted” regression (since it comes in two parts)

k = number of regressors in (each part of the) “unrestricted” regression

3. Perform the test. If the value of the test statistic is greater than the critical value from the F-distribution, which is an $F(k, T-2k)$, then reject the null hypothesis that the parameters are stable over time.

A Chow Test Example

- Consider the following regression for the CAPM β (again) for the returns on Glaxo.
- Say that we are interested in estimating Beta for monthly data from 1981-1992. The model for each sub-period is
- 1981M1 - 1987M10
 $0.24 + 1.2R_{Mt}$ $T = 82$ $RSS_1 = 0.03555$
- 1987M11 - 1992M12
 $0.68 + 1.53R_{Mt}$ $T = 62$ $RSS_2 = 0.00336$
- 1981M1 - 1992M12
 $0.39 + 1.37R_{Mt}$ $T = 144$ $RSS = 0.0434$

A Chow Test Example - Results

- The null hypothesis is

$$H_0: \alpha_1 = \alpha_2 \text{ and } \beta_1 = \beta_2$$

- The unrestricted model is the model where this restriction is not imposed

$$\begin{aligned} \text{Test statistic} &= \frac{0.0434 - (0.0355 + 0.00336)}{0.0355 + 0.00336} \times \frac{144 - 4}{2} \\ &= 7.698 \end{aligned}$$

Compare with 5% $F(2,140) = 3.06$

- We reject H_0 at the 5% level and say that we reject the restriction that the coefficients are the same in the two periods.

The Predictive Failure Test

- Problem with the Chow test is that we need to have enough data to do the regression on both sub-samples, i.e. $T_1 \gg k$, $T_2 \gg k$.
- An alternative formulation is the predictive failure test.
- What we do with the predictive failure test is estimate the regression over a “long” sub-period (i.e. most of the data) and then we predict values for the other period and compare the two.

To calculate the test:

- Run the regression for the whole period (the restricted regression) and obtain the RSS
- Run the regression for the “large” sub-period and obtain the RSS (called RSS_1). Note we call the number of observations T_1 (even though it may come second).

$$\text{Test Statistic} = \frac{RSS - RSS_1}{RSS_1} \times \frac{T_1 - k}{T_2}$$

where T_2 = number of observations we are attempting to “predict”. The test statistic will follow an $F(T_2, T_1 - k)$.

Backwards versus Forwards Predictive Failure Tests

- There are 2 types of predictive failure tests:
 - Forward predictive failure tests, where we keep the last few observations back for forecast testing, e.g. we have observations for 1970Q1-1994Q4. So estimate the model over 1970Q1-1993Q4 and forecast 1994Q1-1994Q4.
 - Backward predictive failure tests, where we attempt to “back-cast” the first few observations, e.g. if we have data for 1970Q1-1994Q4, and we estimate the model over 1971Q1-1994Q4 and backcast 1970Q1-1970Q4.

Predictive Failure Tests – An Example

- We have the following models estimated:

For the CAPM β on Glaxo.

- 1980M1-1991M12

$$0.39 + 1.37R_{Mt} \quad T = 144 \quad RSS = 0.0434$$

- 1980M1-1989M12

$$0.32 + 1.31R_{Mt} \quad T_1 = 120 \quad RSS_1 = 0.0420$$

Can this regression adequately “forecast” the values for the last two years?


$$\text{Test Statistic} = \frac{0.0434 - 0.0420}{0.0420} \times \frac{120 - 2}{24} = 0.164$$

- Compare with $F(24, 118) = 1.66$.

So we do not reject the null hypothesis that the model can adequately predict the last few observations.

How do we decide the sub-parts to use?

- As a rule of thumb, we could use all or some of the following:
 - Plot the dependent variable over time and split the data accordingly to any obvious structural changes in the series, e.g.



- Split the data according to any known important historical events (e.g. stock market crash, new government elected)
- Use all but the last few observations and do a predictive failure test on those.

Measurement Errors

- If there is measurement error in one or more of the explanatory variables, this will violate the assumption that the explanatory variables are non-stochastic
- Sometimes this is also known as the errors-in-variables problem
- Measurement errors can occur in a variety of circumstances, e.g.
 - Macroeconomic variables are almost always estimated quantities (GDP, inflation, and so on), as is most information contained in company accounts
 - Sometimes we cannot observe or obtain data on a variable we require and so we need to use a proxy variable – for instance, many models

Measurement Error in the Explanatory Variable(s)

- Suppose that we wish to estimate a model containing just one explanatory variable, x_t :

- $y_t = \beta_1 + \beta_2 x_t + u_t$,

where u_t is a disturbance term

- Suppose further that x_t is measured with error so that instead of observing its true value, we observe a noisy version, \tilde{x}_t , that comprises the actual x_t plus some additional noise, v_t that is independent of x_t and u_t :

$$\tilde{x}_t = x_t + v_t$$

- Taking the first equation and substituting in for x_t from the second:

$$y_t = \beta_1 + \beta_2(\tilde{x}_t - v_t) + u_t$$

- We can rewrite this equation by separately expressing the composite error term, $(u_t - \beta_2 v_t)$

$$y_t = \beta_1 + \beta_2 \tilde{x}_t + (u_t - \beta_2 v_t)$$

Measurement Error in the Explanatory Variable(s)

- It should be clear from this equation and the one for the explanatory variable measured with error, \tilde{x}_t and the composite error term, $(u_t - \beta_2 v_t)$, are correlated since both depend on v_t
- Thus the requirement that the explanatory variables are non-stochastic does not hold
- This causes the parameters to be estimated inconsistently
- The size of the bias in the estimates will be a function of the variance of the noise in x_t as a proportion of the overall disturbance variance
- If β_2 is positive, the bias will be negative but if β_2 is negative, the bias will be positive
- So the parameter estimate will always be biased towards zero as a result of the measurement noise.

Measurement Error and Tests of the CAPM

- The standard approach to testing the CAPM pioneered by Fama and MacBeth (1973) comprises two stages
- Since the betas are estimated at the first stage rather than being directly observable, they will surely contain measurement error
- The effect of this has sometimes been termed attenuation bias.
- Tests of the CAPM showed that the relationship between beta and returns was smaller than expected, and this is precisely what would happen as a result of measurement error
- Various approaches to solving this issue have been proposed, the most common of which is to use portfolio betas in place of individual betas
- An alternative approach (Shanken, 1992) is to modify the standard errors in the second stage regression to adjust directly for the measurement errors.

Measurement Error in the Explained Variable

- Measurement error in the explained variable is much less serious than in the explanatory variable(s)
- This is one of the motivations for the inclusion of the disturbance term in a regression model
- When the explained variable is measured with error, the disturbance term will in effect be a composite of the usual disturbance term and another source of noise from the measurement error
- Then the parameter estimates will still be consistent and unbiased and the usual formulae for calculating standard errors will still be appropriate
- The only consequence is that the additional noise means the standard errors will be enlarged relative to the situation where there was no measurement error in y .

A Strategy for Building Econometric Models

Our Objective:

- To build a statistically adequate empirical model which
 - satisfies the assumptions of the CLRM
 - is parsimonious
 - has the appropriate theoretical interpretation
 - has the right “shape” - i.e.
 - all signs on coefficients are “correct”
 - all sizes of coefficients are “correct”
 - is capable of explaining the results of all competing models

2 Approaches to Building Econometric Models

- There are 2 popular philosophies of building econometric models: the “specific-to-general” and “general-to-specific” approaches.
- “Specific-to-general” was used almost universally until the mid 1980’s, and involved starting with the simplest model and gradually adding to it.
- Little, if any, diagnostic testing was undertaken. But this meant that all inferences were potentially invalid.
- An alternative and more modern approach to model building is the “LSE” or Hendry “general-to-specific” methodology.
- The advantages of this approach are that it is statistically sensible and also the theory on which the models are based usually has nothing to say about the lag structure of a model.

The General-to-Specific Approach

- First step is to form a “large” model with lots of variables on the right hand side
- This is known as a GUM (generalised unrestricted model)
- At this stage, we want to make sure that the model satisfies all of the assumptions of the CLRM
- If the assumptions are violated, we need to take appropriate actions to remedy this, e.g.
 - taking logs
 - adding lags
 - dummy variables
- We need to do this before testing hypotheses
- Once we have a model which satisfies the assumptions, it could be very big with lots of lags & independent variables

The General-to-Specific Approach: Reparameterising the Model

- The next stage is to reparameterise the model by
 - knocking out very insignificant regressors
 - some coefficients may be insignificantly different from each other, so we can combine them.
- At each stage, we need to check the assumptions are still OK.
- Hopefully at this stage, we have a statistically adequate empirical model which we can use for
 - testing underlying financial theories
 - forecasting future values of the dependent variable
 - formulating policies, etc.

Regression Analysis In Practice - A Further Example: Determinants of Sovereign Credit Ratings

- Cantor and Packer (1996)
Financial background:
- What are sovereign credit ratings and why are we interested in them?
- Two ratings agencies (Moody's and Standard and Poor's) provide credit ratings for many governments.
- Each possible rating is denoted by a grading:

Moody's
Poor's

Standard and

Aaa

AAA

.....

.....

B3

B-

Purposes of the Paper

- to attempt to explain and model how the ratings agencies arrived at their ratings.
- to use the same factors to explain the spreads of sovereign yields above a risk-free proxy
- to determine what factors affect how the sovereign yields react to ratings announcements

Determinants of Sovereign Ratings

- Data

Quantifying the ratings (dependent variable): Aaa/AAA=16, ... , B3/B-=1

- Explanatory variables (units of measurement):

- Per capita income in 1994 (thousands of dollars)
- Average annual GDP growth 1991-1994 (%)
- Average annual inflation 1992-1994 (%)
- Fiscal balance: Average annual government budget surplus as a proportion of GDP 1992-1994 (%)
- External balance: Average annual current account surplus as a proportion of GDP 1992-1994 (%)
- External debt Foreign currency debt as a proportion of exports 1994 (%)
- Dummy for economic development
- Dummy for default history

Income and inflation are transformed to their logarithms.

The model: Linear and estimated using OLS

Explanatory Variable	Expected sign	Dependent Variable			
		Average Rating	Moody's Rating	S&P Rating	Moody's / S&P Difference
Intercept	?	1.442 (0.663)	3.408 (1.379)	-0.524 (-0.223)	3.932** (2.521)
Per capita income	+	1.242*** (5.302)	1.027*** (4.041)	1.458*** (6.048)	-0.431*** (-2.688)
GDP growth	+	0.151 (1.935)	0.130 (1.545)	0.171** (2.132)	-0.040 (0.756)
Inflation	-	-0.611*** (-2.839)	-0.630*** (-2.701)	-0.591*** (2.671)	-0.039 (-0.265)
Fiscal Balance	+	0.073 (1.324)	0.049 (0.818)	0.097* (1.71)	-0.048 (-1.274)
External Balance	+	0.003 (0.314)	0.006 (0.535)	0.001 (0.046)	0.006 (0.779)
External Debt	-	-0.013*** (-5.088)	-0.015*** (-5.365)	-0.011*** (-4.236)	-0.004*** (-2.133)
Development dummy	+	2.776*** (4.25)	2.957*** (4.175)	2.595*** (3.861)	0.362 (0.81)
Default dummy	-	-2.042*** (-3.175)	-1.63** (-2.097)	-2.622*** (-3.962)	1.159*** (2.632)
Adjusted R^2		0.924	0.905	0.926	0.836

Notes: t -ratios in parentheses; *, **, and *** indicate significance at the 10%, 5% and 1% levels respectively. Source: Cantor and Packer (1996). Reprinted with permission from *Institutional Investor*.

Interpreting the Model

From a statistical perspective

- Virtually no diagnostics
- Adjusted R^2 is high
- Look at the residuals: actual rating - fitted rating

From a financial perspective

- Do the coefficients have their expected signs and sizes?

Do Ratings Add to Publicly Available Available Information?

- Now dependent variable is
 - Log (Yield on the sovereign bond - yield on a US treasury bond)

Do Ratings Add to Publicly Available Available Information? Results

Variable	Expected Sign	Dependent Variable: Log (yield spread)		
		(1)	(2)	(3)
Intercept	?	2.105*** (16.148)	0.466 (0.345)	0.074 (0.071)
Average Rating	-	-0.221*** (-19.175)		-0.218*** (-4.276)
Per capita income	-		-0.144 (-0.927)	0.226 (1.523)
GDP growth	-		-0.004 (-0.142)	0.029 (1.227)
Inflation	+		0.108 (1.393)	-0.004 (-0.068)
Fiscal Balance	-		-0.037 (-1.557)	-0.02 (-1.045)
External Balance	-		-0.038 (-1.29)	-0.023 (-1.008)
External Debt	+		0.003*** (2.651)	0.000 (0.095)
Development dummy	-		-0.723*** (-2.059)	-0.38 (-1.341)
Default dummy	+		0.612*** (2.577)	0.085 (0.385)
Adjusted R^2		0.919	0.857	0.914

Notes: t-ratios in parentheses; *, **, and *** indicate significance at the 10%, 5% and 1% levels respectively. Source: Cantor and Packer (1996). Reprinted with permission from *Institutional Investor*.

What Determines How the Market Reacts to Ratings Announcements?

- The sample: Every announcement of a ratings change that occurred between 1987 and 1994 - 79 such announcements spread over 18 countries.
- 39 were actual ratings changes
- 40 were “watchlist / outlook” changes
- The dependent variable: changes in the relative spreads over the US T-bond over a 2-day period at the time of the announcement.

What Determines How the Market Reacts to Ratings Announcements? Explanatory variables.

0 /1 dummies for

- Whether the announcement was positive
- Whether there was an actual ratings change
- Whether the bond was speculative grade
- Whether there had been another ratings announcement in the previous 60 days.

and

- The change in the spread over the previous 60 days.
- The ratings gap between the announcing and the other agency

What Determines How the Market Reacts to Ratings Announcements? Results

Dependent Variable: Log Relative Spread	
Independent variable	Coefficient (<i>t</i> -ratio)
Intercept	-0.02 (-1.4)
Positive announcements	0.01 (0.34)
Ratings changes	-0.01 (-0.37)
Moody's announcements	0.02 (1.51)
Speculative grade	0.03** (2.33)
Change in relative spreads from day -60 to day -1	-0.06 (-1.1)
Rating gap	0.03* (1.7)
Other rating announcements from day -60 to day -1	0.05** (2.15)
Adjusted R^2	0.12

Note: * and ** denote significance at the 10% and 5% levels respectively. Source: Cantor and Packer (1996). Reprinted with permission from *Institutional Investor*.

Conclusions

- 6 factors appear to play a big role in determining sovereign credit ratings - incomes, GDP growth, inflation, external debt, industrialised or not, and default history.
- The ratings provide more information on yields than all of the macro factors put together.
- We cannot determine well what factors influence how the markets will react to ratings announcements.

Comments on the Paper

- Only 49 observations for first set of regressions and 35 for yield regressions and up to 10 regressors
- No attempt at reparameterisation
- Little attempt at diagnostic checking
- Where did the factors (explanatory variables) come from?

Reference and source:

1. Multivariate Time Series Analysis: With R and Financial Applications by Ruey S. Tsay
2. Time Series Analysis by James Douglas Hamilton
3. The Analysis of Time Series: An Introduction with R (Chapman & Hall/CRC Texts in Statistical Science)
4. Machine Learning for Time Series Forecasting with Python by Francesca Lazzeri
5. Time Series Analysis for the Social Sciences (Analytical Methods for Social Research) Part of: Analytical Methods for Social Research (14 Books)
6. Introduction to Probability, Statistics, and Random Processes by Hossein Pishro-Nik
7. Introduction to Time Series and Forecasting (Springer Texts in Statistics) Part of: Springer Texts in Statistics